## Persistent Albuminuria Categories

			A1	A1	A1						
			Normal-Mildly Increased	Moderately Increased	Severely Increased						
			<30 mg/g	30-300 mg/g	>300 mg/g	BMD	NA Intake	Diabetes Treatment			
GFR Categories (ml/min/1.73m²)	Stage 1	Normal or High	≥ 90	1 visit/yr if CKD	1 visit/yr	2 visits/yr		<4 g (<2 g/d if HTN or DM)	Metformin	SGLT-2i	GLP- 1RA
	Stage 2	Mildly Decreased	60-89	1 if CKD	1	2					
	Stage 3a	Mild- Moderately Decreased	45-59	1	2	3					
	Stage 3b	Moderately -Severely Decreased	30-44	2	3	3					
	Stage 4	Severely Decreased	15-29	3	3	4+	Assess for Bone Mineral	<3 g			
	Stage 5	Kidney Failure	<15	4+	4+	4+	Disorder (<2 g/d	(<2 g/d if HTN or DM)			
	Hyper	tension with	out DM	ACE/ARB, CCB, Diuretic	ACE/ARB Suggested	ACE/ARB					
	Hypertensi	ion – with DN	1, GFR > 60	ACE, ARB, CCB, Diuretic	ACE/ARB	ACE/ARB					
	Hypertension – with DM, GFR < 60				ACE/ARB	ACE/ARB					